



## Esperion to Present at the 37th Annual J.P. Morgan Healthcare Conference

January 2, 2019

ANN ARBOR, Michigan, Jan. 02, 2019 (GLOBE NEWSWIRE) -- Esperion (NASDAQ:ESPR), the Lipid Management Company focused on developing and commercializing complementary, convenient, cost-effective, once-daily, oral therapies for the treatment of patients with elevated low density lipoprotein cholesterol (LDL-C), today announced that president and chief executive officer, Tim M. Mayleben, will present at the 37<sup>th</sup> Annual J.P. Morgan Healthcare Conference in San Francisco, CA on Wednesday, January 9, 2019 at 5:30 p.m. Pacific Time / 8:30 p.m. Eastern Time.

A live audio webcast of this event can be accessed on the investor relations section of the Esperion website at [www.esperion.com](http://www.esperion.com). Replay of the webcast will be archived on the Company's website for 90 days following the event.

### Bempedoic Acid / Ezetimibe Combination Pill

Through the complementary mechanisms of action of inhibition of cholesterol synthesis (bempedoic acid) and inhibition of cholesterol absorption (ezetimibe), the bempedoic acid / ezetimibe combination pill is our lead, non-statin, orally available, once-daily, LDL-C lowering therapy. Inhibition of ATP Citrate Lyase (ACL) by bempedoic acid reduces cholesterol biosynthesis and lowers LDL-C by up-regulating the LDL receptor. Inhibition of Niemann-Pick C1-Like 1 (NPC1L1) by ezetimibe results in reduced absorption of cholesterol from the gastrointestinal tract, thereby reducing delivery of cholesterol to the liver, which in turn upregulates the LDL receptors. Phase 3 data demonstrated that this safe and well tolerated combination results in a 35 percent lowering of LDL-C when used with maximally tolerated statins, a 43 percent lowering of LDL-C when used as a monotherapy, and a 34 percent reduction in high sensitivity C-reactive protein (hsCRP).

### Bempedoic Acid

With a targeted mechanism of action, bempedoic acid is a first-in-class, complementary, orally available, once-daily ATP Citrate Lyase (ACL) inhibitor that, reduces cholesterol biosynthesis and lowers LDL-C by up-regulating the LDL receptor. Similar to statins, bempedoic acid also reduces hsCRP, a key marker of inflammation associated with cardiovascular disease. Completed Phase 2 and Phase 3 studies conducted in almost 4,800 patients, and approximately 3,100 patients treated with bempedoic acid, have produced an additional 20 percent LDL-C lowering when used with maximally tolerated statins, up to 30 percent LDL-C lowering as monotherapy, 35 percent LDL-C lowering in combination with ezetimibe when used with maximally tolerated statins, and up to 48 percent LDL-C lowering in combination with ezetimibe as monotherapy.

The effect of bempedoic acid on cardiovascular morbidity and mortality has not yet been determined. The company initiated a global cardiovascular outcomes trial (CVOT) to assess the effects of bempedoic acid on the occurrence of major cardiovascular events in patients with, or at high risk for, cardiovascular disease (CVD) who are only able to tolerate less than the lowest approved daily starting dose of a statin and considered "statin intolerant." The CVOT — known as Cholesterol Lowering via Bempedoic Acid, an ACL-inhibiting Regimen (CLEAR) Outcomes — is an event-driven, global, randomized, double-blind, placebo-controlled study expected to enroll approximately 12,600 patients with hypercholesterolemia and high CVD risk at over 1,000 sites in approximately 30 countries.

### Esperion's Commitment to Patients with Hypercholesterolemia

High levels of LDL-C can lead to a build-up of fat and cholesterol in and on artery walls (known as atherosclerosis), potentially leading to cardiovascular events, including heart attack or stroke. In the U.S., 96 million people, or more than 37 percent of the adult population have elevated LDL-C. There are approximately 18 million people in the U.S. with atherosclerotic cardiovascular disease (ASCVD) who live with elevated levels of LDL-C despite taking maximally tolerated lipid-modifying therapy — including individuals considered statin intolerant — leaving them at high risk for cardiovascular events. More than 50 percent of ASCVD patients who are not able to reach their LDL-C goals with statins alone, need less than a 40 percent reduction to reach their LDL-C threshold.

Esperion's mission as the Lipid Management Company is to deliver once-daily, oral therapies that complement existing oral drugs to provide the additional LDL-C lowering that these patients need.

### The Lipid Management Company

Esperion is the Lipid Management Company passionately committed to developing and commercializing convenient, complementary, cost-effective, once-daily, oral therapies for the treatment of patients with elevated LDL-C. Through scientific and clinical excellence, and a deep understanding of cholesterol biology, the experienced Lipid Management Team at Esperion is committed to developing new LDL-C lowering therapies that will make a substantial impact on reducing global cardiovascular disease; the leading cause of death around the world. Bempedoic acid and the company's lead product candidate, the bempedoic acid / ezetimibe combination pill, are targeted therapies that have been shown to significantly lower elevated LDL-C levels in patients with hypercholesterolemia, including patients inadequately treated with current lipid-modifying therapies. For more information, please visit [www.esperion.com](http://www.esperion.com) and follow us on Twitter at <https://twitter.com/EsperionInc>.

#### Investor Contact:

Alex Schwartz  
Esperion  
734-249-3386  
[aschwartz@esperion.com](mailto:aschwartz@esperion.com)

#### Media Contact:

Elliot Fox  
W2O Group

212-257-6724

[efox@w2ogroup.com](mailto:efox@w2ogroup.com)



Source: Esperion Therapeutics, Inc.